





# **A Future High Statistics Charm Mixing Experiment using the Tevatron**

*Alan Schwartz  
University of Cincinnati*


***First Open Meeting of the SuperKEKB Collaboration  
KEK, Tsukuba, Japan  
11 December 2008***

- ***quick summary of  $D^0$  meson mixing***
- ***a Tevatron experiment: expected  $D^0$  yields***
- ***comparison with B factories/LHCb***
- ***expected sensitivity to CPV***

# $D^0$ meson mixing I:



off-shell ("virtual") states:  $\Delta m$



on-shell states:  $\Delta\Gamma$

| Meson   | f avors    | $\Delta m/\Gamma$ | $\Delta\Gamma/2\Gamma$ | mixing observed? |
|---------|------------|-------------------|------------------------|------------------|
| $K^0$   | $\bar{s}d$ | 0.474             | 0.997                  | yes (1958)       |
| $B^0$   | $\bar{b}d$ | 0.77              | < 1%                   | yes (1987)       |
| $B_s^0$ | $\bar{b}s$ | 27                | $0.15 \pm 0.07$        | yes (2006)       |
| $D^0$   | $cu$       | < 0.029           | $0.011 \pm 0.005$      | yes (2007)       |

- small because:**
- doubly-Cabibbo-suppressed with respect to  $\Gamma_D$
  - GIM mechanism cancellation

# $D^0$ mixing measurements





- **Wrong-sign semileptonic  $D^0(t) \rightarrow K^+ l^- \bar{\nu}$  decays**  
measures  $x^2 + y^2$ , no DCS contamination
- **Wrong-sign hadronic  $D^0(t) \rightarrow K^+ \pi^-$  decays**  
measures  $x' = x \cos \delta + y \sin \delta$ ,  $y' = y \cos \delta - x \sin \delta$ ,  
where  $\delta$  is a strong phase difference
- **Decays to  $CP$  eigenstates:  $D^0(t) \rightarrow K^+ K^-, \pi^+ \pi^-$**   
measures  $y \cos \phi$ , where  $\phi$  is a weak phase difference
- **Wrong-sign hadronic  $D^0 \rightarrow K^+ \pi^- \pi^+ \pi^-, K^+ \pi^- \pi^0$  decays**  
measures  $x^2 + y^2$ ,  $x''$  and  $y''$  with Dalitz plot analysis
- **Dalitz plot analysis of  $D^0(t) \rightarrow K^0 \pi^+ \pi^-$  decays**  
measures  $x, y$
- **Quantum correlations in  $e^+ e^- \rightarrow D^0 \bar{D}^0 (n\pi^0), D^0 \bar{D}^0 \gamma (n\pi^0)$**   
measures  $y, \cos \delta$



# FNAL experiment: yield estimate #1

Scale from HERA-B:  $61.3 \pm 13$   $D^*$ -tagged  $CF D^0 \rightarrow K^- \pi^+$  in  $182 \times 10^6$  hadronic interactions. Multiplying this rate by  $\Gamma(D^0 \rightarrow K^+ \pi^-)/\Gamma(D^0 \rightarrow K^- \pi^+) = 0.377\%$  gives a fractional rate (including (loose) trigger + reconstruction efficiencies) of  $1.3 \times 10^{-9}$  (Reference: I. Abt et al., Eur. Phys. Jour. C52, 531, 2007)



One year of running, assuming 7 MHz interaction rate and trigger efficiency of 50% relative to that of HERA-B:

$$(7 \text{ MHz})(1.4 \times 10^7 \text{ s})(1.3 \times 10^{-9})(0.5) = 64000 \text{ } D^*\text{-tagged } D^0 \rightarrow K^+ \pi^-$$


**192000 in 3 years of running**

# *FNAL experiment: yield estimate #2*


**Scale from E791: 35  $D^*$ -tagged DCS  $D^0 \rightarrow K^+ \pi^-$  in  $5 \times 10^{10}$  hadronic interactions. This gives a fractional rate (including (loose) trigger + reconstruction efficiencies) of  $7 \times 10^{-10}$**  (Reference: E. Aitala et al., PRD 57, 13, 1998)

$p^-$  beam,  
 $p = 500 \text{ GeV}/c$


*CF  $D \rightarrow K^- \pi$*




*CF  $D \rightarrow K^- \pi \pi$*



*DCS  $D \rightarrow K^+ \pi$*



*DCS  $D \rightarrow K^+ \pi \pi$*



**One year of running, assuming 7 MHz interaction rate and the same trigger+reconstruction efficiency as E791:**

$$(7 \text{ MHz})(1.4 \times 10^7 \text{ s})(7 \times 10^{-10}) = 69000 \text{ } D^*\text{-tagged } D^0 \rightarrow K^+ \pi^-$$

**207000 in 3 years of running**



# Comparison with Belle/Babar



**Belle:** 4024  $D^*$ -tagged DCS  $D^0 \rightarrow K^+ \pi^-$  in 400  $\text{fb}^{-1}$  of data  
(Reference: L. Zhang et al., PRL 96, 151801, 2006)



**Babar:** 4030  $D^*$ -tagged DCS  $D^0 \rightarrow K^+ \pi^-$  in 384  $\text{fb}^{-1}$  of data  
(Reference: B. Aubert et al., PRL 98, 211802 (2007))



**Babar has stopped taking data; its final sample is 485  $\text{fb}^{-1}$  (at 4S and continuum)**  
**Belle is now at 839  $\text{fb}^{-1}$ , will probably take another 100  $\text{fb}^{-1}$  at 4S and continuum**



**total sample is 1420  $\text{fb}^{-1}$  or 14600 DCS  $D^0 \rightarrow K^+ \pi^-$  decays,  
8% of the yield of a 3-year Fermilab experiment**

# Comparison with LHCb



**LHCb:** have used Monte Carlo to study sensitivity to  $D^*$ -tagged DCS  $D^0 \rightarrow K^+ \pi^-$  decays

(Reference: P. Spradlin, G. Wilkinson, F. Xing, et al., LHCb public note LHCb-2007-049)

Assuming  $\sigma_{bb} = 500 \mu b$  and estimating several trigger and reconstruction efficiencies, this study concludes that 58000 signal decays will be reconstructed per year ( $2 \text{ fb}^{-1}$  of data)

This is similar to our yield estimate for a Tevatron experiment, **BUT:**

**LHCb must use  $D^*$ 's produced in  $B$  decays, rather than prompt  $D^*$ 's, as the trigger is efficient only for the former. This introduces two issues:**

- (a) the decay time distribution will be a mixture of prompt  $D^*$ 's and  $D^*$ 's from  $B$  decays. These two components will need to be unfolded to measure mixing parameters. (FNAL experiment only has prompt  $D^*$ 's)
- (b) to reconstruct the  $D^*$  vertex position, one must reconstruct a  $B \rightarrow D^* X$  decay. The efficiency for this is estimated to be 51%, and it will add smearing to the  $D^0$  decay time distribution. (FNAL experiment:  $D^*$  vertex is at the target)

# Estimate of sensitivity

Since sample size is very similar, we adopt the results of **LHCb Monte Carlo study** (P. Spradlin et al., Note LHCb-2007-049):

This study found that statistical errors for mixing parameters corresponding to 232500  $D^*$ -tagged  $D^0 \rightarrow K^+\pi^-$  decays,  $S/B = 0.40$ , and  $\sigma_\tau = 75$  ps are:

|               |                       |
|---------------|-----------------------|
| $\delta x'^2$ | $6.4 \times 10^{-5}$  |
| $\delta y'$   | $0.87 \times 10^{-3}$ |

best case after all B factory  
data analyzed:  
(1500 fb<sup>-1</sup>)

|                      |
|----------------------|
| $14 \times 10^{-5}$  |
| $2.2 \times 10^{-3}$ |

This yield corresponds to 3.6 years of running. One also expects a similar improvement in  $y_{CP}$  measured using  $D^0 \rightarrow K^+K^-$  and  $D^0 \rightarrow \pi^+\pi^-$  decays:



|                 |                       |
|-----------------|-----------------------|
| $\delta y_{CP}$ | $0.65 \times 10^{-3}$ |
|-----------------|-----------------------|



# Estimate of sensitivity, cont'd

## Our own toy MC study

(200000  $D^*$ -tagged  $D^0 \rightarrow K^+ \pi^-$  decays,  $S/B = 0.40$ ,  $\sigma_\tau = 75$  ps,  $0.5\tau_D$  cut):



$$\delta x'^2 = 5.8 \times 10^{-5}$$
$$\delta y' = 1.0 \times 10^{-3}$$

(similar to LHCb study)

(RMS's of distribution of residuals  
from 200 toy experiments)

# Estimate of sensitivity, cont'd

## Global fit: fit

24 observables for

8 underlying parameters

$$R_M = \frac{1}{2}(x^2 + y^2)$$

$$2y_{CP} = (|q/p| + |p/q|)y \cos \phi - (|q/p| - |p/q|)x \sin \phi$$

$$2A_\Gamma = (|q/p| - |p/q|)y \cos \phi - (|q/p| + |p/q|)x \sin \phi$$

$$x_{K^0\pi\pi} = x$$

$$y_{K^0\pi\pi} = y$$

$$|q/p|_{K^0\pi\pi} = |q/p|$$

$$\text{Arg}(q/p)_{K^0\pi\pi} = \phi$$

$$\begin{pmatrix} x'' \\ y'' \end{pmatrix}_{K^+\pi^-\pi^0} = \begin{pmatrix} \cos \delta_{K\pi\pi} & \sin \delta_{K\pi\pi} \\ -\sin \delta_{K\pi\pi} & \cos \delta_{K\pi\pi} \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$$

$$\begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{pmatrix} \cos \delta & \sin \delta \\ -\sin \delta & \cos \delta \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$$

$$A_M = \frac{|q/p|^2 - |p/q|^2}{|q/p|^2 + |p/q|^2}$$

$$x'^{\pm} = \left( \frac{1 \pm A_M}{1 \mp A_M} \right)^{1/4} (x' \cos \phi \pm y' \sin \phi)$$

$$y'^{\pm} = \left( \frac{1 \pm A_M}{1 \mp A_M} \right)^{1/4} (y' \cos \phi \mp x' \sin \phi)$$

$$\frac{1}{2} [R(D^0 \rightarrow K^+\pi^-) + \overline{R}(\overline{D}^0 \rightarrow K^-\pi^+)] = R_D$$


$$\frac{R(D^0 \rightarrow K^+\pi^-) - \overline{R}(\overline{D}^0 \rightarrow K^-\pi^+)}{R(D^0 \rightarrow K^+\pi^-) + \overline{R}(\overline{D}^0 \rightarrow K^-\pi^+)} = A_D$$




# Sensitivity to CPV parameters $|q/p|$ , $\phi$

**Global fit:** 24 observables, 8 underlying parameters:

**HFAG now:**




**200000  $D^0 \rightarrow K^+ \pi^-$  decays (Tevatron):**



**CPV is manifest as  $(|q/p|, \phi) \neq (1,0)$  - this would be new physics**

# Summary & comments

- Charm mixing now observed, interest switching to **CP violation** in charm decays.
- Advances since the old Fermilab fix-target experiments make a new experiment much **more powerful/sensitive**. Trigger concepts and prototypes exist (HERA-B, CDF, BTeV, LHCb); pixels and higher rate detectors now developed.
- Notably better sensitivity than all (10 x 2 = 20 years!) B factory data; notably **simpler to analyze** than LHCb data.
- Accelerator and beamline essentially **available**.
- Can significantly improve sensitivity to CPV in the charm system, help **un-tangle whatever signals** appear at the Tevatron or LHC.



# Current Activities

---

**Working group has formed, see:**

<http://www.nevis.columbia.edu/twiki/bin/view/FutureTev/WebHome>

**To subscribe to the listserv:**

- Send an e-mail message to [listserv@fnal.gov](mailto:listserv@fnal.gov)
- Leave the subject line blank
- Type SUBSCRIBE Future\_Charm\_at\_Fermilab FIRSTNAME LASTNAME  
in the body of your message.

**We are now editing a working report  
"Possibilities of a Future Tevatron Program"**



to outline a future physics program at the Tevatron. Such a program would also include a neutrino experiment (J. Conrad) and possibly a hyperon experiment.

***(NOTE: this is not Project X)***